

CHEST[®]

Official publication of the American College of Chest Physicians



Aging and Induced-Sputum Cells


Mario Malerba, Bruno Balbi and Antonio Spanevello

Chest 2005;128:4049-4050
DOI 10.1378/chest.128.6.4049

The online version of this article, along with updated information and services can be found online on the World Wide Web at:
<http://chestjournal.org>

CHEST is the official journal of the American College of Chest Physicians. It has been published monthly since 1935. Copyright 2007 by the American College of Chest Physicians, 3300 Dundee Road, Northbrook IL 60062. All rights reserved. No part of this article or PDF may be reproduced or distributed without the prior written permission of the copyright holder
(<http://www.chestjournal.org/misc/reprints.shtml>). ISSN: 0012-3692.

A M E R I C A N C O L L E G E O F
 C H E S T
P H Y S I C I A N S[®]



communications to the editor

Communications for this section will be published as space and priorities permit. The comments should not exceed 350 words in length, with a maximum of five references; one figure or table can be printed. Exceptions may occur under particular circumstances. Contributions may include comments on articles published in this periodical, or they may be reports of unique educational character. Please submit letters online at <http://mc.manuscriptcentral.com/CHEST>. Please include a cover letter with a complete list of authors (including full first and last names and highest degree), corresponding author's address, phone number, fax number, and e-mail address (if applicable). Specific permission to publish should be cited in the cover letter or appended as a postscript. CHEST reserves the right to edit letters for length and clarity.

Detection of Hypoventilation

To the Editor:

The study by Fu et al (November 2004)¹ documents well that supplemental oxygen impairs the detection of hypoventilation by pulse oximetry, as discussed in the accompanying editorial by Demers.² We regularly perform overnight oximetry on inpatients to screen for sleep apnea/hypopnea and find that the results are very insensitive if performed using supplemental oxygen. Therefore, we perform overnight oximetry either using room air or, if the baseline awake oxygen saturation level is < 90%, using only enough oxygen to bring the awake saturation to approximately 90%.

Fortunately, it is now possible to directly assess hypoventilation using continuous transcutaneous carbon dioxide tension monitoring.³ We now routinely monitor transcutaneous carbon dioxide pressure in patients who are at high risk for hypoventilation in our ventilator weaning program. We find monitoring to be very helpful during the initial tracheostomy using a mask or during overnight periods when the patient is not receiving ventilation, as well as during bronchoscopies. The advantages of monitoring cutaneous carbon dioxide tension over monitoring with end-tidal carbon dioxide tension, which we also use, include allowing continuous measurement, not requiring deep exhalation, and making accurate measurements in patients with high dead space ventilation.

The device (CO-OXSYS Monitor, SenTec AG; Therwil, Switzerland) has been used in Europe^{4,5} and is now available in the United States (Aspen Medical Products Inc; Irvine, CA). The monitor allows us to closely follow transcutaneous carbon dioxide pressure using a small probe that clips on the ear lobe. It usually takes about 5 min to equilibrate and then tracks carbon dioxide pressure closely, along with oxygen saturation.

Douglas C. Johnson, MD
Boston, MA

Reproduction of this article is prohibited without written permission from the American College of Chest Physicians (www.chestjournal.org/misc/reprints.shtml).

Correspondence to: Douglas C. Johnson, MD, Spaulding Reha-

bilitation Hospital, 125 Nashua St, Boston, MA 02114; e-mail: djohnson5@partners.org

REFERENCES

- 1 Fu ES, Downs JB, Schweiger JW, Miguel RV, et al. Supplemental oxygen impairs detection of hypoventilation by pulse oximetry. *Chest* 2004; 126:1552-1558
- 2 Demers B. The oximeter: boon or bane? *Chest* 2004; 126:1399-1401
- 3 Chhajed PN, Heuss LT, Tamm M. Cutaneous carbon dioxide tension monitoring in adults. *Curr Opin Anaesthesiol* 2004; 17:521-525
- 4 Heuss LT, Chhajed PN, Schnieper P, Hirt T, et al. Combined pulse oximetry/cutaneous carbon dioxide tension monitoring during colonoscopies: pilot study with a smart ear clip. *Digestion* 2004; 70:152-158
- 5 Chhajed PN, Rajasekaran R, Kaegi B, et al. Cutaneous carbon dioxide tension monitoring might enhance patient safety during flexible bronchoscopy and medical thoracoscopy [abstract]. *Chest* 2004; 126(suppl):822S

Aging and Induced-Sputum Cells

To the Editor:

We read with interest the article by Thomas et al¹ (December 2004) on the influence of age on induced sputum in normal subjects. We think that the discussion on the possible physiopathologic mechanisms could be deepened. Although an influence of advancing age on lung cellularity in healthy subjects has been already described,² reference values for cell counts in induced sputum in healthy adults > 50 years old are not available. The possible explanations of the results found by Thomas et al¹ could be an impairment in humoral lung immunity in older healthy subjects compared with younger healthy subjects,³ and the presence of a low-grade inflammation in the lower respiratory tracts of many asymptomatic, older subjects.⁴ In particular, a previous study² has reported in BAL fluid of the older healthy individuals an increase in CD4+/CD8+ lymphocytes ratio probably due to a repeated antigenic stimulation or irritation by environmental substances of the immune cells in the lower respiratory tract during the years. The recurrent antigen stimulation on the immune cells in the lung could be demonstrated by the decreasing with age of CD19+ B lymphocytes that represent the B cells not yet differentiated into antibody-secreting cells, suggesting that B cells on mucosal surface of airways in older subjects has been driven to differentiate by previous repeated antigen stimulations. The low-grade inflammation in the airways observed in older subjects might be related to the decline in the lung function that starts in the fourth to fifth decade of life in normal never-smoker humans. The mechanism by which neutrophils are recruited to within the airways in older healthy subjects is still unclear. A number of neutrophil chemoattractants can be secreted by inflammatory cells that reside in the airways, and epithelial cells can release cytokines, such as IL-8, which have a potent chemoattractant activity for neutrophils.⁵ Low-grade persistent inflammation may occur because of the loss of factors that normally down-regulate the inflammatory response to pollutants

or repetitive antigenic stimulations, combined with advancing age. Epithelial cells could be a significant source of neutrophil chemoattractants, which contributes to a low-grade inflammation in older subjects. Persistent, low-grade inflammation could damage elastin and perhaps lead to the age-associated loss of elastin fibers. Therefore, considering that many patients affected by asthma or COPD who increasingly perform induced sputum are often > 50 years old, these findings deserve further investigations.

Mario Malerba, MD
University of Brescia
Bruno Balbi, MD
Istituto di Gussago Brescia
Brescia, Italy
Antonio Spanevello, MD
Istituto di Cassano Murge
Bari, Italy

Reproduction of this article is prohibited without written permission from the American College of Chest Physicians (www.chestjournal.org/misc/reprints.shtml).

Correspondence to: Mario Malerba, MD, Department of Internal Medicine, University of Brescia, I Medicina, Spedali Civili di Brescia, Piazza Spedali Civili n 1, Brescia, Italy; e-mail: malerba@master.cci.unibs.it

REFERENCES

- 1 Thomas RA, Green RH, Brightling CE, et al. The influence of age on induced sputum differential cell counts in normal subjects. *Chest* 2004; 126:1811–1814
- 2 Meyer KC, Ershler W, Rosenthal NS, et al. Immune dysregulation in the aging human lung. *Am J Respir Crit Care Med* 1996; 153:1072–1079
- 3 Ammann AJ, Schiffman G, Austrian R. The antibody responses to pneumococcal capsular polysaccharides in aged individuals. *Proc Soc Exp Biol Med* 1980; 164:312–316
- 4 Thompson AB, Scholer SG, Daughton DM, et al. Altered epithelial lining fluid parameters in old normal individuals. *J Gerontol* 1992; 47:M171–M176
- 5 Abramson SL, Malech HL, Gallin JI. Neutrophils. In: Crystal RG, West JB, ed. *The lung: scientific foundations*. New York, NY: Raven Press, 1991, 553–563

Ribavirin Should Be Tested in Clinical Trials in Combination With Other Antiviral Agents for Severe Acute Respiratory Syndrome

To the Editor:

We read with interest the article in *CHEST* by Chiou et al (July 2005)¹ and offer the following comments. The ribavirin-treated patients had higher lactate dehydrogenase levels, a well-known adverse prognostic factor in severe acute respiratory syndrome (SARS). The nonsignificantly higher mortality could be due to the more severe disease in this group. Viral load, another important predictor of mortality, was not available.² Moreover, Figure 1 seemed inaccurate: the survival in ribavirin-treated patients should be 0.88 at day 30 (5 of 44 patients died) instead of 0.71.

Classifying the ribavirin-treated patients into hypoxemic and nonhypoxemic subgroups (Table 2) and attributing the higher mortality in the hypoxemic subgroup to ribavirin was problematic, as both subgroups were treated with an identical protocol of ribavirin. From the data presented, a more likely explanation for

the more severe drop in hemoglobin in the hypoxemic subgroup was that they had more severe disease. The survival curves in Figure 4 also appeared inaccurate: the survival in patients with drop in hemoglobin > 2 g/dL should be 0.69 (5 of 16 patients died) instead of 0.45. Hence, the result of the log-rank test ($p = 0.007$) needs to be justified.

Only factors that were potentially associated with hypoxemia were analyzed in Table 2. No univariate or multivariate analyses on factors related to death were reported. The conclusion that hemoglobin level was the only factor associated with death was not supported by the data presented.

In Figure 6, the shaded triangles were supposed to represent the hemoglobin of patients who were hypoxemic and had received ribavirin. There were 22 triangles, but there should only be 17 patients. In addition, expressing the survival of individual patients by proportion (y-axis) is difficult to understand.

Therefore, there is no convincing evidence that ribavirin has contributed to a life-threatening drop in hemoglobin or mortality in this report. As of today, three independent studies^{3–5} have shown ribavirin to have *in vitro* activities against SARS-coronavirus, alone or in combination with other agents. Ribavirin should be tested in future randomized controlled studies in combination with other potential antiviral agents for SARS.

REFERENCES

- 1 Chiou HE, Liu CL, Buttrey MJ, et al. Adverse effects of ribavirin and outcome in severe acute respiratory syndrome: experience in two medical centers. *Chest* 2005; 128:263–272
- 2 Chu CM, Poon LL, Cheng VC, et al. Initial viral load and the outcomes of severe acute respiratory syndrome (SARS). *Can Med Assoc J* 2004; 171:1349–1352
- 3 Chu CM, Cheng VC, Hung IF, et al. Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. *Thorax* 2004; 59:252–256
- 4 Chen F, Chan KH, Jiang Y, et al. *In vitro* susceptibility of 10 clinical isolates of SARS coronavirus to selected antiviral compounds. *J Clin Virol* 2004; 31:69–75
- 5 Morgenstern B, Michaelis M, Baer PC, et al. Ribavirin and interferon- β synergistically inhibit SARS-associated coronavirus replication in animal and human cell lines. *Biochem Biophys Res Commun* 2005; 326:905–908

Chung-Ming Chu, MD, FCCP
Kin-Sang Chan, MBBS, FCCP
United Christian Hospital
Hong Kong

Reproduction of this article is prohibited without written permission from the American College of Chest Physicians (www.chestjournal.org/misc/reprints.shtml).

Correspondence to: Chung-Ming Chu, MD, FCCP, United Christian Hospital, Hong Kong, PRC

A Modified Percutaneous Tracheostomy Technique Without Bronchoscopic Guidance

A Note of Concern

To the Editor:

We read with interest the article in *CHEST* by Paran and colleagues (September 2004)¹ on a modified percutaneous tra-

Aging and Induced-Sputum Cells
Mario Malerba, Bruno Balbi and Antonio Spanevello
Chest 2005;128;4049-4050
DOI 10.1378/chest.128.6.4049

This information is current as of July 21, 2008

Updated Information & Services	Updated information and services, including high-resolution figures, can be found at: http://chestjournal.org/cgi/content/full/128/6/4049
References	This article cites 4 articles, 2 of which you can access for free at: http://chestjournal.org/cgi/content/full/128/6/4049#BIBL
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://chestjournal.org/misc/reprints.shtml
Reprints	Information about ordering reprints can be found online: http://chestjournal.org/misc/reprints.shtml
Email alerting service	Receive free email alerts when new articles cite this article sign up in the box at the top right corner of the online article.
Images in PowerPoint format	Figures that appear in CHEST articles can be downloaded for teaching purposes in PowerPoint slide format. See any online article figure for directions.

